

AD-A051 179

FLORIDA STATE UNIV TALLAHASSEE DEPT OF STATISTICS  
A RELIABILITY GROWTH MODEL INVOLVING DEPENDENT COMPONENTS.(U)

F/G 12/1

OCT 77 N LANGBERG, F PROSCHAN

AFOSR-74-2581

UNCLASSIFIED

FSU-STATISTICS-M437

AFOSR-TR-78-0275

NL

| OF |

AD  
A051179



END  
DATE  
FILMED  
4 -78  
DDC

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

1. REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER		
4. TITLE (and Subtitle)	5. TYPE OF REPORT PERIOD COVERED		
6. AUTHOR(s)	7. PERFORMING ORG. REPORT NUMBER		
8. CONTRACT OR GRANT NUMBER(s)	9. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS		
10. CONTROLLING OFFICE NAME AND ADDRESS	11. REPORT DATE		
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	13. NUMBER OF PAGES		
14. DISTRIBUTION STATEMENT (of this Report)	15. SECURITY CLASS. (of this report)		
15a. DECLASSIFICATION/DOWNGRADING SCHEDULE			
16. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
17. SUPPLEMENTARY NOTES			
18. KEY WORDS (Continue on reverse side if necessary and identify by block number)			
19. ABSTRACT (Continue on reverse side if necessary and identify by block number)			

DD FORM 1473 EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

AD A051179  
DDC FILE COPY

400 277

JOB

next Page

20. Abstract

the limiting distributions in the sequence of dependent models "correctly" correspond to the limiting distributions in the sequence of independent models.

These results have applications in reliability growth models and in biomedical competing risk models in which the competing risks are increasing with age; in these models dependency is permitted among the random variables.

ACCESSION for	
NTIS	Write Section <input checked="" type="checkbox"/>
DDC	BRI Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
DI	SP CIAL
A	

2

A RELIABILITY GROWTH MODEL INVOLVING  
DEPENDENT COMPONENTS


by

N. Langberg and F. Proschan<sup>1</sup>

FSU Statistics Report M437  
AFOSR Technical Report No. 77

October, 1977

The Florida State University  
Department of Statistics  
Tallahassee, Florida 32306



<sup>1</sup>Research sponsored by the Air Force Office of Scientific Research, AFSC, USAF, under Grant AFOSR-74-2581D. The United States Government is authorized to reproduce and distribute reprints for governmental purposes.

Approved for public release  
distribution unlimited.



A RELIABILITY GROWTH MODEL INVOLVING  
DEPENDENT COMPONENTS

by

N. Langberg and F. Proschan<sup>1</sup>

Abstract

Earlier papers have shown how to convert competing risk models involving dependent random variables into models involving only independent random variables, while simultaneously preserving the distribution of the minimum and the probabilities of the various failure patterns. In the present paper, we consider a sequence of such conversions occurring at successive points in chronological time in which the independent random variables are becoming stochastically larger. We obtain results which essentially demonstrate that the limiting distributions in the sequence of dependent models "correctly" correspond to the limiting distributions in the sequence of independent models.

These results have applications in reliability growth models and in biomedical competing risk models in which the competing risks are increasing with age; in these models dependency is permitted among the random variables.

<sup>1</sup>Research sponsored by the Air Force Office of Scientific Research, AFSC, USAF, under Grant AFOSR-74-2581D. The United States Government is authorized to reproduce and distribute reprints for governmental purposes.

# A RELIABILITY GROWTH MODEL INVOLVING DEPENDENT COMPONENTS

by

N. Langberg and F. Proschan  
Florida State University

1. Introduction. In several earlier papers, Tsiatis (1975), Miller (1977), and Langberg, Proschan, and Quinzi (1978) show how to convert various competing risk models involving dependent random variables into models involving only independent random variables, while simultaneously preserving the distribution of the minimum and the probabilities corresponding to certain "failure patterns". Explicit equations are presented which yield the distributions of the independent variables. In a more recent paper, Langberg, Proschan, and Quinzi (1977) develop statistical estimators of parameters of interest in competing risk models in which causes or times of death (in the biomedical context) or of failure (in the reliability context) are not necessarily independent.

In the present paper we present a result which should prove to be basic in converting reliability growth models involving dependent failure times into equivalent models involving only independent random variables. Analogous applications exist in the biomedical field in which survival functions may be decreasing,<sup>†</sup> rather than increasing, with chronological time. The probabilistic theorem presented will be useful in inference in reliability growth models involving dependent failure times.

---

<sup>†</sup>Throughout the paper we use "decreasing" in place of "nonincreasing" and "increasing" in place of "nondecreasing".

2. The Reliability Growth Model. The notation and terminology are as in Langberg, Proschan, and Quinzi (1978).

Consider a series system of  $n$  components undergoing improvement as time passes. At fixed chronological time  $u$ ,  $0 \leq u < u_0$  ( $u_0$  possibly infinite) component  $i$  has random lifelength  $T_i(u)$  for  $i = 1, \dots, n$ , where  $T_1(u), \dots, T_n(u)$  are not necessarily mutually independent. We say that failure pattern  $I$  occurs if the simultaneous failures of the components in subset  $I$  of  $\{1, \dots, n\}$  and of no other components causes (i.e., coincides with) the failure of the system. Define

$$\xi(\underline{T}(u)) = \begin{cases} I & \text{if failure pattern } I \text{ occurs at time } u \\ \emptyset & \text{otherwise.} \end{cases}$$

Let  $\underline{S}(u)$  and  $\underline{T}(u)$  represent the vectors of component life lengths of two systems whose system life lengths are  $S(u)$  and  $T(u)$ , respectively. We say that the two systems are equivalent in life lengths and patterns at time  $u$

$(\underline{S}(u) \stackrel{LP}{=} \underline{T}(u))$  if  $P(S(u) > t, \xi(\underline{S}(u)) = I) = P(T(u) > t, \xi(\underline{T}(u)) = I)$  for each  $t \geq 0$  and each  $I \subset \{1, \dots, n\}$ . Thus, two systems which are equivalent in life length and patterns are such that (i) their life lengths have the same distribution and (ii) the corresponding failure patterns in the two systems have the same probability of occurrence.

The main result of Langberg, Proschan, and Quinzi (1978) may be restated in the context of our model as follows:

2.1. Theorem. For fixed  $u$ ,  $0 \leq u < u_0$ , let  $T(u) = \min(T_i(u), 1 \leq i \leq n)$  denote the life length of an  $n$ -component series system, where  $T_i(u)$  represents the life length of component  $i$ ,  $i = 1, \dots, n$ . Define  $\bar{F}_{u,I}(t) = P(T(u) > t, \xi(\underline{T}(u)) = I)$ ,  $F_{u,I}(t) = P(T(u) \leq t, \xi(\underline{T}(u)) = I)$ ,  $\bar{F}_u(t) = P(T(u) > t)$ , and



$\alpha(F_u) = \sup\{x: \bar{F}_u(x) > 0\}$ . Then the following statements hold:

(i) A necessary and sufficient condition for the existence of a set of independent random variables  $(H_I, I \in \{1, \dots, n\})$  which satisfy  $H(u) \stackrel{LP}{=} T(u)$ , where  $H(u) = \min(H_I(u), I \in \{1, \dots, n\})$ , is that the sets of discontinuities of the  $F_{u,I}$  be pairwise disjoint.

(ii) The distributions of  $H_I(u), I \in \{1, \dots, n\}$ , are uniquely determined on the interval  $[0, \alpha(F_u))$  as follows:

$$\bar{G}_{u,I}(t) \equiv P(H_I(u) > t) = \exp\left[-\int_0^t (dF_{u,I}^C / \bar{F}_u)\right]. \quad (2.1)$$

$$\cdot \prod_{a_{u,I}(I,j)} \bar{F}_u(a_{u,I}(I,j)) / [\bar{F}_u(a_{u,I}(I,j)) + f_{u,I}(a_{u,I}(I,j))],$$

$0 \leq t < \alpha(F_u)$ , where  $F_{u,I}^C$  is the continuous part of  $F_{u,I}$ ,  $\{a_{u,I}(I,j)\}$  is the set of discontinuities of  $F_{u,I}$ , in  $[0, t]$  and  $f_{u,I}(a_{u,I}(I,j))$  is the size of the jumps of  $F_{u,I}$  at  $a_{u,I}(I,j)$ .

(Note that  $G_{u,I}$  may place mass at infinity.)

The survival probabilities  $\bar{G}_{u,I} \equiv 1 - G_{u,I}$  for  $H_I(u)$ , the time until a shock occurs which simultaneously destroys subset  $I$ , for  $I \in \{1, \dots, n\}$  are obtained by solving the identities:

$$\prod_{I \in \{1, \dots, n\}} \bar{G}_{u,I}(t) = \bar{F}_u(t), \quad t \geq 0, \quad (2.2)$$

and

$$\int_t^\infty \prod_{J \neq I} \bar{G}_{u,J} dG_{u,I} = \bar{F}_{u,I}(t), \quad t \geq 0. \quad (2.3)$$



Thus at each fixed instant  $u$  of chronological time, we may "replace" the original series system of dependent components by a set of mutually independent sources of shock; a given source of shock fails a corresponding subset of components. Moreover, the replacement is so chosen as to yield the same joint distribution of failure pattern and system life length as possessed by the original system. Finally, the distribution of the time of each type of shock occurrence is explicitly specified in (2.1).

Suppose now that the system is experiencing reliability growth; specifically, assume that:

(i)  $\bar{G}_{u,I}$  is increasing in  $u$ ,  $0 \leq u < u_0$  for each  $I \in \{1, \dots, n\}$ , so that  $H_I(u)$  is stochastically increasing in  $u$ ,  $0 \leq u < u_0$  for each  $I \in \{1, \dots, n\}$ . That is, each type of shock is occurring with decreasing frequency as chronological time passes. This, in turn, implies that the system life length  $T(u)$  is stochastically increasing in  $u$ . Since  $\bar{G}_{u,I}(t)$  is monotone increasing in  $u < u_0$ , it follows immediately that the  $\bar{G}_{u,I}(t)$  converge to  $\inf_{x>t} \sup_{u<u_0} \bar{G}_{u,I}(x)$  for each  $t \geq 0$ ,  $I \in \{1, \dots, n\}$ ; call this limit  $\bar{G}_I(t)$ .  $\bar{G}_I(t)$  is  $^0$  a survival function, with mass possibly at infinity.

Three basic questions now arise:

- (1) Does the joint distribution  $F_{u,I}$  of failure pattern and life length converge to a joint distribution as  $u \rightarrow u_0$  for each  $I \in \{1, \dots, n\}$ ?
- (2) If such limiting joint distributions exist, do they coincide respectively with the solutions (call them  $F_I$ ) of (2.1) corresponding to the limit distributions  $G_I$ ,  $I \in \{1, \dots, n\}$  (as we would hope)?
- (3) Does the system lifelength distribution  $F_u$  converge to a lifelength distribution as  $u \rightarrow u_0$ ; if so, does it coincide with the solution  $F$  obtained from (2.2) where the subscript  $u$  is omitted?

In the next section we prove that under mild, reasonable conditions, affirmative answers exist for all three questions.

3. Limit Results in the Reliability Growth Model. In this section we give convenient sufficient conditions to yield affirmative answers to questions (1), (2), and (3) at the end of Section 2, and present proofs of our results.

We are considering the model of Section 2. We make the following two additional assumptions:

(ii) The sets of discontinuities of the  $G_{u,I}$  are pairwise disjoint for each fixed time point  $u$ ,  $0 \leq u < u_0$ .

(iii) The sets of discontinuities of the  $G_I$  are pairwise disjoint.

The main result may now be stated:

3.1. Theorem. Assume the reliability growth model specified in Section 2 and assume (i), (ii), and (iii). Then

(1)  $\lim_{u \rightarrow u_0} F_{u,I}(t)$  exists for each  $t \geq 0$  and each  $I \in \{1, \dots, n\}$ ,

(2)  $\lim_{u \rightarrow u_0} F_{u,I}(t) = F_I(t)$  for each  $t \geq 0$  and each  $I \in \{1, \dots, n\}$ ,

(3)  $\lim_{u \rightarrow u_0} F_u(t)$  exists and  $= F(t)$  for each  $t \geq 0$ ; moreover,  $\bar{F}_u(t)$  is increasing upward to  $\bar{F}(t)$  as  $u$  increases upward to  $u_0$ .

To prove Theorem 3.1, we will make use of

3.2 Lemma. Let  $u_k$  be an increasing sequence converging to  $u_0$ . Then for each  $t \geq 0$  and  $I \in \{1, \dots, n\}$ ,  $\limsup_{k \rightarrow \infty} \bar{F}_{u_k,I}(t) \leq \bar{F}_I(t)$ .

Proof of Lemma. By (2.3), we may write:

$$\bar{F}_{u_k,I}(t) = \int_t^\infty \prod_{J \neq I} \bar{G}_{u_k,J} dG_{u_k,I} \leq \int_t^\infty \prod_{J \neq I} \bar{G}_J dG_{u_k,I},$$

since each  $\bar{G}_{u_k,J} \leq \bar{G}_J$ . Next note that

$$\begin{aligned} \int_t^\infty \prod_{J \neq I} \bar{G}_J dG_{u, I} &= \int_t^\infty [\bar{G}_{u_k, I}(t) - \bar{G}_{u_k, I}(x^-)] d_x [1 - \prod_{J \neq I} \bar{G}_J(x)] \\ &\leq \int_t^\infty [\bar{G}_{u_k, I}(t) - \bar{G}_{u_k, I}(x)] d_x [1 - \prod_{J \neq I} \bar{G}_J(x)]. \end{aligned}$$

But by the Lebesgue dominated convergence theorem,

$$\begin{aligned} \lim_{k \rightarrow \infty} \int_t^\infty [\bar{G}_{u_k, I}(t) - \bar{G}_{u_k, I}(x)] d_x [1 - \prod_{J \neq I} \bar{G}_J(x)] \\ = \int_t^\infty [\bar{G}_I(t) - \bar{G}_I(x)] d_x [1 - \prod_{J \neq I} \bar{G}_J(x)] \\ = \int_t^\infty \prod_{J \neq I} \bar{G}_J(x) dG_I(x) = \bar{F}_I(t), \end{aligned}$$

by assumption (iii). The desired conclusion now follows. ||

We may now prove Theorem 3.1.

Proof of Theorem 3.1 (1), (2). Let  $\{u_k\}$ ,  $0 \leq u_k < u_0$ , be an arbitrary sequence converging to  $u_0$  and let  $t \geq 0$ . By (2.2),

$$\sum_{I \in \{1, \dots, n\}} \bar{F}_{u_k, I}(t) = \prod_{I \in \{1, \dots, n\}} \bar{G}_{u_k, I}(t) =$$

$$\limsup_{k \rightarrow \infty} \prod_{I \in \{1, \dots, n\}} \bar{G}_{u_k, I}(t) = \limsup_{k \rightarrow \infty} \sum_{I \in \{1, \dots, n\}} \bar{F}_{u_k, I}(t).$$

But

$$\limsup_{k \rightarrow \infty} \sum_{I \in \{1, \dots, n\}} \bar{F}_{u_k, I}(t) \leq \sum_{I \in \{1, \dots, n\}} \limsup_{k \rightarrow \infty} \bar{F}_{u_k, I}(t) \leq \sum_{I \in \{1, \dots, n\}} \bar{F}_I(t),$$

by Lemma 3.2. Now by assumption (iii) and the appropriate version of (2.2) we have:

$$\sum_{I \in \{1, \dots, n\}} \bar{F}_I(t) = \prod_{I \in \{1, \dots, n\}} \bar{G}_I(t).$$



Recalling that  $\bar{G}_I(t) = \lim_{k \rightarrow \infty} \bar{G}_{u_k, I}(t)$  from Section 2 and summarizing the inequalities above, we may state:

$$\prod_{I \in \{1, \dots, n\}} \bar{G}_I(t) \leq \sum_{I \in \{1, \dots, n\}} \limsup_{k \rightarrow \infty} \bar{F}_{u_k, I}(t) \leq \sum_{I \in \{1, \dots, n\}} \bar{F}_I(t) = \prod_{I \in \{1, \dots, n\}} \bar{G}_I(t).$$

The desired conclusions (1) and (2) now follow immediately.

Proof of Theorem 3.1 (3). By (2.2),  $\lim_{u \rightarrow u_0} \bar{F}_u(t)$  exists since each  $\lim_{u \rightarrow u_0} \bar{G}_{u, I}$  exists. Next note that

$$\begin{aligned} \bar{F}(t) &= \sum_{I \in \{1, \dots, n\}} \bar{F}_I(t) = \prod_{I \in \{1, \dots, n\}} \bar{G}_I(t) = \lim_{u \rightarrow u_0} \prod_{I \in \{1, \dots, n\}} \bar{G}_{u, I}(t) = \\ &= \lim_{u \rightarrow u_0} \sum_{I \in \{1, \dots, n\}} \bar{F}_{u, I}(t) = \lim_{u \rightarrow u_0} \bar{F}_u(t). \end{aligned}$$

The monotonicity of  $\bar{F}_u(t)$  in  $u$  is a consequence of the monotonicity of each  $\bar{G}_{u, I}$ . ||

#### 4. Extensions, Modifications, and Generalizations.

4.1. Remark. In the reliability growth model discussed above, we assumed that reliability growth was occurring during the interval of chronological time  $[0, t_0]$ . Obviously, the conclusions of Theorem 3.1 may be obtained by assuming that reliability growth occurs at the time points of any subset  $U$  of  $[0, \infty)$  (or, for that matter, over the entire half-line). As examples, consider:

(a)  $U = \{u, 2u, 3u, \dots\}$ , where  $u > 0$ ,

(b)  $U = \{0 \leq u_1 < u_2 < u_3 < \dots\}$

(c)  $U = \{[0, u_0] < u_1 < u_2 < \dots\}$ ,

etc. Moreover, the set  $U$  need not be deterministic, but may be random.

4.2. Remark. In the model of Section 2 and the corresponding result, Theorem 3.1, we assumed that shock intervals from each source were stochastically increasing with chronological time. It is clear that by reversing inequalities and the direction of monotonicity, we can obtain a dual to Theorem 3.1 in which the intervals between shocks from each source are stochastically decreasing with chronological time. System reliability (or in the biomedical context, organism survival probability) would decrease with chronological time.

4.3. Remark. The model of Section 2 and the limit results of Section 3 were formulated in terms of reliability growth. An equally important and useful application exists in the context of competing risks of death of a biological organism from a variety of diseases, accidents, or other causes. In this context, Remark 4.2 is especially relevant. We consider a situation in which an organism is subject to death as a result of any of a number of diseases or combinations of diseases. As time passes, the organism becomes more

vulnerable to each of the diseases. Note that the model now appropriate differs from the model of Section 2 only in the direction of the monotonicity; i.e., the different types of shocks are becoming stochastically more frequent, rather than, as in the first model, stochastically less frequent.

4.4. Remark. In the reliability growth model, we are able to deduce monotonicity of  $\bar{F}_u$ , system reliability as a function of chronological time  $u$ . However it is not necessarily true that each  $\bar{F}_{u,I}$  is monotone in  $u$ . This is a consequence of the fact that as some modes of failure become less likely, others may become more likely.

4.5. Remark. Assumption (iii) of Theorem 3.1 can not be obtained as a consequence of assumptions (i) and (ii). A simple counterexample is available to verify this assertion.



References

- Langberg, N., Proschan, F., and Quinzi, A. J. (1977). Estimation of dependent life lengths, with application to the theory of competing risks. (In preparation.)
- Langberg, N., Proschan, F., and Quinzi, A. J. (1978). Converting dependent models into independent ones, preserving essential features. Ann. Prob. 6 (in press)
- Miller, D. R. (1977). A note on independence of multivariate lifetimes in competing risk models. Ann. Statist. 5 576-579.
- Tsiatis, A. (1975). A nonidentifiability aspect of the problem of competing risks. Proc. Nat. Acad. Sci. USA. 72 20-22.

UNCLASSIFIED  
SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1. REPORT NUMBER FSU No. M437 AFOSR No. 77	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE A Reliability Growth Model Involving Dependent Variables	5. TYPE OF REPORT & PERIOD COVERED Technical Report	
	6. PERFORMING ORG. REPORT NUMBER FSU Statistics Report M437	
7. AUTHOR(s) N. Langberg and F. Proschan	8. CONTRACT OR GRANT NUMBER(s) AFOSR-74-2581D	
9. PERFORMING ORGANIZATION NAME & ADDRESS The Florida State University Department of Statistics Tallahassee, Florida 32306	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
11. CONTROLLING OFFICE NAME & ADDRESS The U.S. Air Force Air Force Office of Scientific Research Bolling Air Force Base, D.C. 20332	12. REPORT DATE October, 1977	
	13. NUMBER OF PAGES 10	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	15. SECURITY CLASS (of this report) Unclassified	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS competing risks, reliability growth, dependent components, limit theorems, failure patterns, system failure, equivalent models.		
20. ABSTRACT		

Earlier papers have shown how to convert competing risk models involving dependent random variables into models involving only independent random variables, while simultaneously preserving the distribution of the minimum and the probabilities of the various failure patterns. In the present paper, we consider a sequence of such conversions occurring at successive points in chronological time in which the independent random variables are becoming stochastically larger. We obtain results which essentially demonstrate that the limiting distributions in the sequence of dependent models "correctly" corresponds to the limiting distributions in the sequence of independent models.

These results have applications in reliability growth models and in biomedical competing risk models in which the competing risks are increasing with age; in these models dependency is permitted among the random variables.